

LÖWNER CHAINS WITH COMPLEX LEADING COEFFICIENT

IKKEI HOTTA

ABSTRACT. In this paper we confirm that several crucial theorems due to Pommerenke and Becker for the theory of Löwner chains work well without normalization on the complex-valued first coefficient. As applications of those considerations, some new univalent and quasiconformal extension criteria are given in the last section.

1. Introduction

Let $\mathbb{C}$ and $\hat{\mathbb{C}}$ denote the complex plane and the Riemann sphere respectively, and let $\mathbb{D}_r = \{z \in \mathbb{C} : |z| < r \leq 1\}$ and $\mathbb{D} = \mathbb{D}_1$. We denote by $\mathcal{A}$ the class of functions $f(z)$ normalized so that $f(0) = f'(0) - 1 = 0$ which are analytic on $\mathbb{D}$ and $\mathcal{S}$ the subclass of $\mathcal{A}$ whose members are univalent on $\mathbb{D}$.

Let $f_t(z) = f(z, t) = \sum_{n=1}^{\infty} a_n(t)z^n$, $a_1(t) \neq 0$, be a function defined on $\mathbb{D} \times [0, \infty)$ and analytic in $\mathbb{D}$ for each $t \in [0, \infty)$, where $a_1(t)$ is a complex-valued, locally absolutely continuous function on $[0, \infty)$ and $\lim_{t \rightarrow \infty} |a_1(t)| = \infty$. Then $f_t(z)$ is called a *Löwner chain* if $f_t(z)$ satisfies the following conditions;

1. $f_t(z)$ is univalent in $\mathbb{D}$ for each $t \in [0, \infty)$,
2. $f_s(\mathbb{D}) \subsetneq f_t(\mathbb{D})$ for $0 \leq s < t < \infty$.

If $a_1(t) = e^t$, then we say that $f_t(z)$ is a *standard Löwner chain*.

It is known that if $f_t(z)$ is a Löwner chain then $f_{t_n}(\mathbb{D}) \rightarrow f_{t_0}(\mathbb{D})$ if $t_n \rightarrow t_0 \in [0, \infty)$ and $f_{t_n}(\mathbb{D}) \rightarrow \mathbb{C}$ if $t_n \rightarrow \infty$ in the sense of kernel convergence with respect to the origin. However, the converse is not true in general (see for instance [6, pp.136–138] and [8, pp.94–97]).

Löwner chains and several related theorems serve as a powerful tool for the theory of univalent functions. In those researches, it seems that many authors deal with the case when the first coefficient $a_1(t)$ of a Löwner chain is a real-valued function. Mostly, they only require the condition $a_1(t) \neq 0$ and it is not clear that $a_1(t)$ is either real- or complex-valued. Certainly, the fact is known that $a_1(t)$ can be taken to be complex-valued (e.g. [2]) and actually it is used by some authors as a tool (e.g. [14], [16], [9]). However, usually there is no explicit mention that the first coefficient of Löwner chains is admitted to be a complex-valued function.

Date: Oct. 26, 2009.

2000 Mathematics Subject Classification. Primary 30C55, 30C80, Secondary 30C45, 30C62.

Key words and phrases. Löwner(Loewner) chain, quasiconformal mapping, univalent function.

One of our purposes is to give a precise statement and a proof to our key theorems for Löwner chains with complex first coefficient, and to confirm that those theorems work well without normalization. In consequence, the obscurity which is mentioned above will be completely removed.

A Löwner chain without normalization on the first derivative has some new applications to univalence and quasiconformal extension criteria. Another purpose of this paper is to derive those new criteria as a benefit of our considerations. We discuss it in the last section.

2. Related theorems for Löwner chains

The following necessary and sufficient condition for a standard Löwner chain is known;

Theorem A ([14],[15]). *Let $0 < r_0 \leq 1$. Let $h(z, t) = e^t z + \sum_{n=2}^{\infty} c_n(t) z^n$ be a function defined on $\mathbb{D} \times [0, \infty)$ and analytic in $\mathbb{D}$ for each $t \in [0, \infty)$. Then the function $h(z, t)$ is a standard Löwner chain if and only if the following two conditions are satisfied;*

- i) *The function $h(z, t)$ is analytic in $z \in \mathbb{D}_{r_0}$ for each $t \in [0, \infty)$, absolutely continuous in $t \in [0, \infty)$ for each $z \in \mathbb{D}_{r_0}$ and satisfies*

$$|h(z, t)| \leq K_0 e^t \quad (z \in \mathbb{D}_{r_0}, t \in [0, \infty)) \quad (1)$$

for some positive constants K_0 .

- ii) *There exists a function $p(z, t)$ analytic in $z \in \mathbb{D}$ for each $t \in [0, \infty)$ and measurable in $t \in [0, \infty)$ for each $z \in \mathbb{D}$ satisfying*

$$\operatorname{Re} p(z, t) > 0 \quad (z \in \mathbb{D}, t \in [0, \infty))$$

such that

$$\dot{h}(z, t) = zh'(z, t)p(z, t) \quad (z \in \mathbb{D}_{r_0}, \text{ a.e. } t \in [0, \infty)) \quad (2)$$

where $\dot{h} = \partial h / \partial t$ and $h' = \partial h / \partial z$.

We will show first that Theorem A can be generalized for a Löwner chain which has the complex-valued first coefficient as the following form;

Theorem A'. *Let $0 < r_1 \leq 1$. Let $f(z, t) = \sum_{n=1}^{\infty} a_n(t) z^n$, $a_1(t) \neq 0$, be a function defined on $\mathbb{D} \times [0, \infty)$ and analytic in $\mathbb{D}$ for each $t \in [0, \infty)$, where $a_1(t)$ is a complex-valued, locally absolutely continuous function on $[0, \infty)$ and $\lim_{t \rightarrow \infty} |a_1(t)| = \infty$. Then the function $f(z, t)$ is a Löwner chain if and only if the following conditions are satisfied;*

- i') The function $f(z, t)$ is analytic in $\mathbb{D}_{r_1}$ for each $t \in [0, \infty)$, locally absolutely continuous in $[0, \infty)$ for each $z \in \mathbb{D}_{r_1}$ and

$$|f(z, t)| \leq K_1 |a_1(t)| \quad (z \in \mathbb{D}_{r_1}, \text{ a.e. } t \in [0, \infty)) \quad (3)$$

for some positive constants K_1 .

- ii') There exists a function $p(z, t)$ analytic in $\mathbb{D}$ for each $t \in [0, \infty)$ and measurable in $[0, \infty)$ for each $z \in \mathbb{D}$ satisfying

$$\operatorname{Re} p(z, t) > 0 \quad (z \in \mathbb{D}, t \in [0, \infty))$$

such that

$$\dot{f}(z, t) = z f'(z, t) p(z, t) \quad (z \in \mathbb{D}_{r_1}, \text{ a.e. } t \in [0, \infty)) \quad (4)$$

where $\dot{f} = \partial f / \partial t$ and $f' = \partial f / \partial z$.

Proof. Let $f(z, t) = \sum_{n=1}^{\infty} a_n(t) z^n$ be a Löwner chain, where $a_1(t) \neq 0$ is a complex-valued locally absolutely continuous function on $[0, \infty)$.

First we set $\lambda(t) := -\arg a_1(t)$ which is locally absolutely continuous with respect to t and define

$$g(z, t) = \sum_{n=1}^{\infty} b_n(t) z^n := f(e^{i\lambda(t)} z, t). \quad (5)$$

This yields

$$\frac{\dot{g}(z, t)}{z g'(z, t)} = \frac{\dot{f}(e^{i\lambda(t)} z, t)}{e^{i\lambda(t)} z f'(e^{i\lambda(t)} z, t)} + i\lambda'(t).$$

It follows that $\operatorname{Re} \dot{g}(z, t) / z g'(z, t) > 0$ if and only if $\operatorname{Re} \dot{f}(z, t) / z f'(z, t) > 0$ for all $z \in \mathbb{D}$ and almost all $t \in [0, \infty)$. We remark that $b_1(t) = |a_1(t)|$ is a strictly increasing positive function in $t \in [0, \infty)$, and the inverse function b_1^{-1} is defined in $[|a_1(0)|, \infty)$ and maps this interval onto $[0, \infty)$.

Next, let

$$h(z, t) := \frac{1}{|a_1(0)|} g(z, b_1^{-1}(|a_1(0)| e^t)). \quad (6)$$

It follows immediately from these reparametrizations that $f(z, t)$ is a Löwner chain if and only if $h(z, t)$ is a standard Löwner chain. We note that therefore h , namely b_1^{-1} also, is absolutely continuous with respect to t by Theorem A.

We also have

$$\frac{\dot{h}(z, t)}{z h'(z, t)} = \frac{\dot{g}(z, b_1^{-1}(|a_1(0)| e^t))}{z g'(z, b_1^{-1}(|a_1(0)| e^t))} \cdot (b_1^{-1}(|a_1(0)| e^t))'.$$

from (6). Since $(b_1^{-1}(t))' > 0$ for almost all $t \in [0, \infty)$, $\operatorname{Re} \dot{h}(z, t) / z h'(z, t) > 0$ if and only if $\operatorname{Re} \dot{g}(z, t) / z g'(z, t) > 0$ for all $z \in \mathbb{D}$ and $t \in [0, \infty)$. We can see easily that $h(z, t)$ satisfies the condition (1) if and only if $f(z, t)$ satisfies (3) with $K_1 = K_0 / |a_1(0)|^2$, and the other properties of the sufficient part of Theorem A are preserved by the reparametrizations (5) and (6) with $r_0 = r_1$. Consequently, all the necessary and sufficient conditions of Theorem A' follows from Theorem A. $\square$

We remark that a similar argument about normalization of Löwner chains as above can be found in [2]. In fact, $p(z, t)$ in (4) is normalized by

$$p^*(e^{i\beta(t)}z, \alpha(t)) = \frac{1}{\operatorname{Re} p(0, t)} [p(z, t) - i\operatorname{Im} p(0, t)]$$

where

$$\alpha(t) = \int_0^t \operatorname{Re} p(0, \tau) d\tau, \quad \beta(t) = \int_0^t \operatorname{Im} p(0, \tau) d\tau,$$

which is equivalent to $f'(0, t) = \exp\{\alpha(t) + i\beta(t)\}$.

Theorem A appears in [12, p.4] without proof, and the similar change of variables as (5) and (6) is in [8, p.95]. However, it is not clear that whether $a_1(t) = f'(0, t)$ can be taken to be complex-valued or not.

The following theorem which is essentially same as Theorem A' is often used to show univalence for an analytic function, apart from the theory of Löwner chains;

Theorem B ([14]). *Let $0 < r_0 \leq 1$. Let $f(z, t) = a_1(t)z + \sum_{n=2}^{\infty} a_n(t)z^n$, $a_1(t) \neq 0$, be analytic for each $t \in [0, \infty)$ in $\mathbb{D}_{r_0}$ and locally absolutely continuous in $[0, \infty)$, locally uniformly with respect to $\mathbb{D}_{r_0}$, where $a_1(t)$ is a complex-valued function on $[0, \infty)$. For almost all $t \in [0, \infty)$ suppose*

$$\dot{f}(z, t) = z f'(z, t) p(z, t) \quad (z \in \mathbb{D}_{r_0}, t \in [0, \infty))$$

where $p(z, t)$ is analytic in $\mathbb{D}$ and satisfies $\operatorname{Re} p(z, t) > 0$, $z \in \mathbb{D}$. If $|a_1(t)| \rightarrow \infty$ for $t \rightarrow \infty$ and if $\{f(z, t)/a_1(t)\}$ forms a normal family in $\mathbb{D}_{r_0}$, then for each $t \in [0, \infty)$ $f(z, t)$ can be continued analytically in $\mathbb{D}$ and gives a univalent function.

Proof. By using the previous reparametrization argument, it is enough to think about the case when $f(z, t)$ is a standard Löwner chain. Following the lines of the proof of [14, Folgerung 3] one can obtain our assertion. $\square$

Furthermore, we shall look at the next theorem which is due to Becker. Here, a sense-preserving homeomorphism f of $G \subset \mathbb{C}$ is called k -quasiconformal if f_z and $f_{\bar{z}}$, the partial derivatives in z and $\bar{z}$ in the distributional sense, are locally integrable on G and satisfy $|f_{\bar{z}}| \leq k|f_z|$ almost everywhere in G , where $k \in [0, 1)$.

Theorem C ([1]). *Suppose that $h_t(z) = h(z, t)$ is a standard Löwner chain for which $p(z, t)$ in (2) satisfies the condition*

$$\begin{aligned} p(z, t) \in U(k) &:= \left\{ w \in \mathbb{C} : \left| \frac{w-1}{w+1} \right| \leq k \right\} \\ &= \left\{ w \in \mathbb{C} : \left| w - \frac{1+k^2}{1-k^2} \right| \leq \frac{2k}{1-k^2} \right\} \end{aligned}$$

for all $z \in \mathbb{D}$ and $t \in [0, \infty)$. Then $h(z, t)$ admits a continuous extension to $\overline{\mathbb{D}}$ for each $t \geq 0$ and the map $\hat{h}$ defined by

$$\hat{h}(z) = \begin{cases} h(z, 0), & \text{if } |z| < 1, \\ h\left(\frac{z}{|z|}, \log |z|\right), & \text{if } |z| \geq 1, \end{cases} \quad (7)$$

is a k -quasiconformal extension of h_0 to $\mathbb{C}$.

For the proof, see e.g. [3]. The above theorem is also generalized for a Löwner chain $f(z, t)$ with the complex first coefficient;

Theorem C' ([2]). Suppose that $f_t(z) = f(z, t)$ is a Löwner chain for which $p(z, t)$ in (2) satisfies the condition $p(z, t) \in U(k)$ for all $z \in \mathbb{D}$ and $t \in [0, \infty)$, where $a_1(t) = f'(0, t)$ is a complex-valued function on $[0, \infty)$. Then $f_t(z)$ admits a continuous extension to $\overline{\mathbb{D}}$ for each $t \geq 0$ and the map $\hat{f}$ given in (7) is a k -quasiconformal extension of f_0 to $\mathbb{C}$.

For the proof, see [2]. Indeed, it is enough to repeat the original proof in [3] with Theorem A' for $f(z, t)$.

3. Applications

In this section we shall show some univalence and quasiconformal extension criteria as a consequence of the considerations in the previous section.

3.1. Convex combinations. A function $f \in \mathcal{A}$ is called *convex* if f is univalent and $f(\mathbb{D})$ is a convex domain. It is well known that $f \in \mathcal{A}$ is convex if and only if f satisfies $\operatorname{Re} \{1 + zf''(z)/f'(z)\} > 0$ for all $z \in \mathbb{D}$. The author showed in [10] that if f is convex then $\alpha f(z) + (1 - \alpha)zf'(z)$ with $\alpha \in [0, 1]$ is univalent in $\mathbb{D}$. This result is extended as following with the aid of Theorem A';

Theorem 1. Let α be a complex number with $|2\alpha - 1| \leq 1$. If $f \in \mathcal{A}$ is a convex function, then the function

$$\alpha f(z) + (1 - \alpha)zf'(z) \quad (8)$$

is univalent in $\mathbb{D}$.

Proof. Let

$$f_t(z) = \alpha f(z) + e^t(1 - \alpha)zf'(z).$$

Then $\lim_{t \rightarrow \infty} |f'_t(0)| = \lim_{t \rightarrow \infty} |\alpha + (1 - \alpha)e^t| = \infty$. Furthermore

$$\frac{1}{p(z, t)} = \frac{zf'_t(z)}{\dot{f}_t(z)} = \frac{1}{e^t} \left(\frac{\alpha}{1 - \alpha} \right) + 1 + \frac{zf''(z)}{f'(z)}.$$

Since $\operatorname{Re} \alpha/(1 - \alpha) > 0$, it turns out that $f_t(z)$ is a Löwner chain by the assumption and Theorem A'. In particular f_0 is univalent in $\mathbb{D}$ which is our assertion. $\square$

It can be shown that the function (8) is close-to-convex by using convolution technique (Li-Mei Wang, personal communications). On the other hand, by using Löwner's method we can give a simple proof for univalence of (8).

3.2. Spirallike functions. A function $f \in \mathcal{A}$ is called α -spirallike and known to be univalent if f satisfies

$$\operatorname{Re} \left\{ e^{i\alpha} \frac{zf'(z)}{f(z)} \right\} > 0$$

for a real number $\alpha \in (-\pi/2, \pi/2)$ in $\mathbb{D}$. If $\alpha = 0$ then we say that f is *starlike*. It is known [15] that the standard Löwner chain

$$h_t(z) = e^{(1-ia)t} f(e^{iat}z) \quad (9)$$

with $a = \tan \alpha$ corresponds to an α -spirallike function because it follows from calculations that

$$p(z, t) = \frac{\dot{h}_t(z)}{zh'_t(z)} = ia + \frac{1}{\cos \alpha} \left(e^{-i\alpha} \frac{f(e^{iat}z)}{e^{iat}zf'(e^{iat}z)} \right) \quad (10)$$

and therefore $\operatorname{Re} p(z, t) > 0$ implies α -spirallikeness of f . If we apply Theorem A and Theorem C to the standard Löwner chain (9), then we obtain the next proposition. Here, let us denote that $U(\alpha, k)$ the hyperbolic disk in the tilted half plane $\{z \in \mathbb{C} : \operatorname{Re} e^{i\alpha}z > 0\}$ centered at 1 with radius $\operatorname{arctanh} k$, $0 \leq k < 1$, i.e.,

$$U(\alpha, k) = \left\{ w \in \mathbb{C} : \left| w - \frac{1 + e^{-2i\alpha}k^2}{1 - k^2} \right| \leq \frac{2k \cos \alpha}{1 - k^2} \right\}.$$

It is clear that $U(0, k) = U(k)$.

Proposition 2. *Let $\alpha \in (-\pi/2, \pi/2)$ and $k \in [0, 1)$. For $f \in \mathcal{A}$, if*

$$\frac{zf'(z)}{f(z)} \in U(\alpha, k)$$

for all $z \in \mathbb{D}$, then f has a k -quasiconformal extension to $\mathbb{C}$.

The case when $\alpha = 0$ appears e.g. in [5].

Proof. Let $w = zf'(z)/f(z)$. Then applying Theorem A and Theorem C to (10), it can be deduced that if w satisfies

$$\left| \frac{1}{w} - \frac{1 + e^{2i\alpha}k^2}{1 - k^2} \right| \leq \frac{2k \cos \alpha}{1 - k^2}$$

then, f has a k -quasiconformal extension to $\mathbb{C}$. The inequality implies $1/w \in U(-\alpha, k)$ which is equivalent to $w \in U(\alpha, k)$ $\square$

On the other hand, by constructing a Löwner chain without normalization on the first derivative and applying Theorem A' and Theorem C' we have the following another result;

Theorem 3. Let $\alpha \in (-\pi/2, \pi/2)$ and $k \in [|\tan(\alpha/2)|, 1)$. For $f \in \mathcal{A}$, if

$$e^{i\alpha} \frac{zf'(z)}{f(z)} \in U(k)$$

for all $z \in \mathbb{D}$, then f has a k -quasiconformal extension to $\mathbb{C}$.

Proof. Let c be a complex constant with $\operatorname{Re} c > 0$. If we set

$$f_t(z) = e^{ct} f(z),$$

then $\lim_{t \rightarrow \infty} |f'_t(0)| = \lim_{t \rightarrow \infty} |e^{ct}| = \infty$ since $\operatorname{Re} c > 0$. A calculation shows that

$$\frac{1}{p(z, t)} = \frac{1}{c} \frac{zf'(z)}{f(z)}.$$

Therefore we obtain our theorem if we put $c = e^{-i\alpha}$ and apply Theorem A' and Theorem C'. $\square$

In [11], a quasiconformal extension criterion for the class of strongly spirallike functions has been studied. It shows that if there exists a $\beta \in [0, 1)$ such that $e^{i\alpha} zf'(z)/f(z)$ lies in the sectoral domain $\{z : |\arg z| < \pi\beta/2\}$ for all $z \in \mathbb{D}$, then f can be extended to a k -quasiconformal automorphism of $\widehat{\mathbb{C}}$. The dilatation of the extended mapping is implicitly given by $k = \sin(\pi\beta/2)$. The same result was proved also in [7, Korollar 5.3] though it does not mention the dilatation k . A special case of Theorem 3 when $\alpha = 0$ was provided in [4] as a consequence of an extension of Theorem C.

3.3. Bazilevič functions 1. Let $\alpha > 0$ and $\beta \in \mathbb{R}$. In the following two subsections we deal with two quasiconformal extension criteria for a Bazilevič function of type (α, β) . Here, A function $f \in \mathcal{A}$ is called *Bazilevič of type (α, β)* if

$$f(z) = \left[(\alpha + i\beta) \int_0^z g(\zeta)^\alpha h(\zeta) \zeta^{i\beta-1} d\zeta \right]^{1/(\alpha+i\beta)} \quad (11)$$

for a starlike univalent function $g \in \mathcal{A}$ and an analytic function h with $h(0) = 1$ satisfying $\operatorname{Re}(e^{i\lambda} h) > 0$ in $\mathbb{D}$ for some $\lambda \in \mathbb{R}$. We denote by $\mathcal{B}(\alpha, \beta)$ the class of such functions.

If the above h in (11) satisfies $|\arg e^{i\lambda} h| \leq \pi\gamma/2$ in $\mathbb{D}$ for some $\lambda \in [0, 1]$ ($0 \leq \gamma \leq 1$ with $h(0) = 1$), then we say that f is a Bazilevič function of order γ , and denote by $f \in \mathcal{B}(\alpha, \beta, \gamma)$. This notion appears in [7]* which is devoted to the study of the class $\mathcal{B}(\alpha, \beta, \gamma)$ and contains the following quasiconformal extension criterion; *Let $\alpha > 0, \beta \in \mathbb{R}$ and $\gamma \in [0, 1)$. Then for $f \in \mathcal{B}(\alpha, \beta, \gamma)$ the following two conditions imply that $f(\partial\mathbb{D})$ is a rectifiable quasicircle, in particular, f has a quasiconformal extension to $\widehat{\mathbb{C}}$;*

*The author would like to thank Professor Wolfram Koepf for his help under which the author could obtain a copy of the dissertation.

1. $\alpha < \frac{1-\gamma}{4}$,
2. There exists a starlike function $g \in \mathcal{A}$ in (11) such that

$$\lim_{r \rightarrow 1} \left[\frac{\log \max_{|z|=r} |g(z)|}{\log \frac{1}{1-r}} \right] < \frac{1-\gamma}{2\alpha}. \quad (12)$$

It is known that the limit value of the left-hand side in (12) exists for all starlike functions in $\mathcal{A}$ (see e.g. [13]). The above theorem does not estimate the dilatation of the extended quasiconformal mapping explicitly.

It follows from a result of Sheil-Small [17] that

$$\operatorname{Re} \left\{ 1 + \frac{zf''(z)}{f'(z)} + (\alpha + i\beta - 1) \frac{zf'(z)}{f(z)} \right\} > 0$$

for all $z \in \mathbb{D}$ implies that $f \in \mathcal{B}(\alpha, \beta)$. Now we refine this result to k -quasiconformal extension criterion (compare with [9, Theorem 2]);

Theorem 4. Let $\alpha > 0$, $\beta \in \mathbb{R}$, $k \in [0, 1)$ and $f \in \mathcal{A}$. If f satisfies

$$\left[1 + \frac{zf''(z)}{f'(z)} + (\alpha + i\beta - 1) \frac{zf'(z)}{f(z)} \right] \in U(k) \quad (13)$$

for all $z \in \mathbb{D}$, then f can be extended to a k -quasiconformal automorphism of $\mathbb{C}$.

Proof. Let

$$f_t(z) = f(z) \left\{ 1 + (e^t - 1) \frac{zf'(z)}{f(z)} \right\}^{1/(\alpha + i\beta)}.$$

In that case $a_1(t) = f'_t(0) = e^{t/(\alpha + i\beta)}$ and therefore $\lim_{t \rightarrow \infty} |a_1(t)| = \infty$ since $\alpha > 0$. A straightforward calculation shows that

$$\frac{1}{p(z, t)} = \frac{1}{e^t} (\alpha + i\beta) + \left(1 - \frac{1}{e^t} \right) \left(1 + \frac{zf''(z)}{f'(z)} + (\alpha + i\beta - 1) \frac{zf'(z)}{f(z)} \right).$$

Since the assumption (13) implies $\alpha + i\beta \in U(k)$ (consider the case $z = 0$), we have $1/p(z, t) \in U(k)$ by (13). Consequently, it follows from Theorem A' and Theorem C' that f has a k -quasiconformal extension to $\mathbb{C}$. $\square$

3.4. Bazilevič functions 2. A Löwner chain for a Bazilevič function is known [14] as

$$f_t(z) = \left[(\alpha + i\beta) \int_0^z g(\zeta)^\alpha \left\{ h(\zeta) + th_0(\zeta) \right\} \zeta^{i\beta-1} d\zeta \right]^{1/(\alpha + i\beta)} \quad (14)$$

where

$$h_0(z) = i\beta + \alpha \frac{zg'(z)}{g(z)}.$$

But we cannot apply Theorem C to (14) because

$$\frac{1}{p(z, t)} = h(z) + th_0(z)$$

and hence it does not satisfy the assumption (7) when t tends to ∞ . We can avoid this obstacle by reparametrizing t with $e^t - 1$;

Theorem 5. *Let $\alpha > 0$, $\beta \in \mathbb{R}$ and $k \in [0, 1)$. For $f \in \mathcal{B}(\alpha, \beta)$, we suppose that functions h and g of (11) satisfy $h(z) \in U(k)$ and $i\beta + \alpha(zg'(z)/g(z)) \in U(k)$ for all $z \in \mathbb{D}$, respectively. Then f can be extended to a k -quasiconformal automorphism of $\mathbb{C}$.*

Proof. Let

$$f(z, t) = \left[(\alpha + i\beta) \int_0^z g(\zeta)^\alpha [h(\zeta) + (e^t - 1)h_0(\zeta)] \zeta^{i\beta-1} d\zeta \right]^{1/(\alpha+i\beta)}$$

where

$$h_0(z) = i\beta + \alpha \frac{zg'(z)}{g(z)}.$$

Since

$$\dot{f}_t(z) = f(z, t)^{1-(\alpha+i\beta)} \cdot e^t g(z)^\alpha z^{i\beta}$$

and

$$zf'_t(z) = f(z, t)^{1-(\alpha+i\beta)} \cdot g(z)^\alpha [h(z) + (e^t - 1)h_0(z)] z^{i\beta},$$

we have

$$\frac{1}{p(z, t)} = \frac{1}{e^t} h(z) + \left(1 - \frac{1}{e^t}\right) h_0(z).$$

We also have

$$f'_t(0) = (1 + (e^t - 1)(\alpha + i\beta))^{1/(\alpha+i\beta)}$$

and therefore $\lim_{t \rightarrow \infty} |f'_t(0)| = \infty$. Consequently, our assertion follows from Theorem A' and Theorem C'. $\square$

ACKNOWLEDGEMENT

The author is deeply grateful to Professor Toshiyuki Sugawa for many valuable suggestions and comments that have improved this paper. The author also thanks the referee for reading the manuscript carefully and giving several helpful comments.

REFERENCES

1. J. Becker, *Löwnersche Differentialgleichung und quasikonform fortsetzbare schlichte Funktionen*, J. Reine Angew. Math. **255** (1972), 23–43.
2. ———, *Über die Lösungsstruktur einer Differentialgleichung in der konformen Abbildung*, J. Reine Angew. Math. **285** (1976), 66–74.
3. ———, *Conformal mappings with quasiconformal extensions*, Aspects of contemporary complex analysis, Academic Press, London, 1980, pp. 37–77.
4. Th. Betker, *Löwner chains and quasiconformal extensions*, Complex Variables Theory Appl. **20** (1992), no. 1-4, 107–111.

5. J. E. Brown, *Quasiconformal extensions for some geometric subclasses of univalent functions*, Internat. J. Math. Math. Sci. **7** (1984), no. 1, 187–195.
6. John B. Conway, *Functions of one complex variable. II*, Graduate Texts in Mathematics, vol. 159, Springer-Verlag, New York, 1995.
7. U. Gall, *Über das Randverhalten von Bazilevič-Funktionen*, Dissertation an der Technischen Universität Berlin, 1986.
8. I. Graham and G. Kohr, *Geometric function theory in one and higher dimensions*, Monographs and Textbooks in Pure and Applied Mathematics, vol. 255, New York, 2003.
9. I. Hotta, *Ruscheweyh's univalent criterion and quasiconformal extensions*, preprint (arxiv:0910.4744).
10. ———, *Explicit quasiconformal extensions and Löwner chains*, Proc. Japan Acad. Ser. A Math. Sci. **85** (2009), no. 8, 108–111.
11. J. G. Krzyż, *Quasiconformal extensions of some special univalent functions*, Colloq. Math. **51** (1987), 189–193.
12. S. S. Miller and P. T. Mocanu, *Differential subordinations, theory and applications*, Monographs and Textbooks in Pure and Applied Mathematics, vol. 225, New York, 2000.
13. Ch. Pommerenke, *On starlike and convex functions*, J. London Math. Soc. **37** (1962), 209–224.
14. ———, *Über die Subordination analytischer Funktionen*, J. Reine Angew. Math. **218** (1965), 159–173.
15. ———, *Univalent functions*, Vandenhoeck & Ruprecht, Göttingen, 1975.
16. S. Ruscheweyh, *An extension of Becker's univalence condition*, Math. Ann. **220** (1976), no. 3, 285–290.
17. T. Sheil-Small, *On Bazilevič functions*, Quart. J. Math. Oxford Ser. (2) **23** (1972), 135–142.

DIVISION OF MATHEMATICS, GRADUATE SCHOOL OF INFORMATION SCIENCES, TOHOKU UNIVERSITY, 6-3-09 ARAMAKI-AZA-AOBA, AOBA-KU, SENDAI 980-8579, JAPAN, PHONE : +81-22-795-4636
E-mail address: ikkeihotta@ims.is.tohoku.ac.jp
URL: <http://www.ims.is.tohoku.ac.jp/~ikkeihotta>